

QUALITY EVALUATION OF *STEVIA REBAUDIANA* CULTIVATED IN FARMERS FIELD

Chauhan Nirpendra, Lohani Hema, Andola Harish Chandra*

Centre for Aromatic Plants, Industrial Estate, Selaqui-248197 Dehradun, Uttarakhand, India

Received on: 01/09/11 Revised on: 03/10/11 Accepted on: 20/10/11

*Corresponding author

Email: andolah@rediffmail.com

ABSTRACT

Among the potential industrial crops *Stevia rebaudiana* introduced in newly developed state Uttarakhand for cultivation as well as quality evaluations of farmers produced and its comparative assessment with other parts of northern areas. *Stevia rebaudiana* is a natural, non-caloric, sweet-tasting plant used globally for its sweeter properties. *Stevia rebaudiana* cultivated in different region of north India, stevioside vary 6.0- 9.5%, rebaudioside vary ranging 1.60-3.90 % within different location.

KEY WORDS *Stevia rebaudiana*, low calorie sweetener, stevioside, glycosides, cultivated Farmers fields.

INTRODUCTION

Stevia rebaudiana (Bert.) is an herbaceous perennial plants of the Asteraceae family it is native to praguay. Leaves of plant produced zero- caloric diterpene glycosides (stevioside and rebaudioside), a non – calorie nutritive, high, potency sweetener and substitute to sucrose (Introducing *Stevia rebaudiana* a natural zero caloric Sweetener)¹ Eight diterpene glycoside with sweetening properties have been identified in leaf tissues of *stevia*. The four Major glycosides are stevioside, rebaudioside A, rebaudioside C, dulcoside was report² The Asin market consumed over 85 % of the global supply of fluffy white crystalline stevia extract. The current extract market is 1.5 million kg processed 12 million kg of the stevia leaf. *Stevia* leaf³. Stevioside is one of the active constituents which are available abundantly in the leaf of the stevia (5-10 %) on dry weight basis. *Stevia* acts as favour enhancer it is hypoglycemic contraceptive, cardiovascular and antimicrobial activity it is also used to weight loss digestive and skin problems⁴. The leaf extract of these plants has been used traditionally in the treatment of diabetes. The crop was first established in Japan. Extremely used by food and flavors industry, worldwide demand for high potency sweeteners is increasing and especially with blending of different sweeteners, the demand for alternative is expected to increase. The sweet herb *Stevia rebaudiana* Bertoni produces an alternative sweetener with the added advantage that *Stevia* sweeteners are natural plant products. *Stevia* sweeteners have functional and sensory properties superior to those of many other high potency sweeteners. *Stevia* contains stevioside, a glycoside that is 200-350 times sweeter than sucrose⁵. It is useful for people with diabetes, hypoglycemia, and candidiasis. The sweetness in *Stevia* is mainly attributed to two compounds i.e. stevioside and rebaudioside⁶. Environmental factors including biotic and abiotic stimuli carbon –nutrient balance, genotype, and ontogenesis usually control and regulate the biosynthesis of secondary metabolites. It is well reported phenolic and flavonoids greatly affect by UV radiation and habitats conditions⁷. Present Communication deals the quality evaluations of stevioside and rebaudioside in *Stevia rebaudiana* cultivated in farmers fields.

MATERIALS AND METHODS

Plant Material Collection

Sample of *Stevia rebaudiana* collected from ten different locations namely Viz Vikasnagar, Sahaspur, Dehardun, Herbartpur, Jaspur, Haldwani, Pantnagar, Meeruth, Panjab,(Ludiana), and Lucknow of *Stevia rebaudiana* cultivated in different Parts of North India.

Extraction of Plant Material

Sample of stevie sweeteners (100 mg) were dissolved in 100 ml volumetric flask with HPLC grade water 95% ethyl alcohol. The

solutions were passed through a C8 Cartridge (Waters USA) prior to HPLC analysis.

Linearity Response

Five concentrations of stevioside and rebaudioside with range 0.3 to 1.6mg/ml ranges were prepared by dissolving and diluting in mobile phase. The linear response of stevioside and rebaudioside 0.997, 0.999 respectively

HPLC Condition

The samples were analyzed using HPLC system (Waters USA 600 pump) having carbohydrate column (250 x 4.6 mm, x 5 μ) analytical column. Mobile phase ratio 80:20 Acetonitrile: Water), flow rate 1 ml/min, run time 15 minutes and detector wavelength was set at 210 nm. Standard stevioside and rebaudioside were procured from Sigma Chemicals USA, and other chemicals from Merck Co, Mumbai India.

RESULTS

Result of present study show in (Table-1), study revealed that sample collected from Vikasnagar, contain (9.57 %) greater amount of stevioside and sample collected from Lucknow possessed (6.0%) lowest amount of stevioside in case of rebaudioside Vikasnagar sample contain maximum amount (3.90 %), similarly Lucknow locality contain minimum amount of rebaudioside (3.80 %).

DISCUSSION

The present study showed the greater variation in glycosides content different region of north India. It is well reported that vegetative growth is reduced when temperature is below 200C and day length to 16 hours and increasing light intensity can enhance vegetative growth and stevioside level⁶. However the chemical components and their contents of crude drug often vary depending on the geographical locations of the habitats seasons, plant parts, processing procedure also harvesting time including various factors, such as an age of plant season, microbial attack grazing, radiation, competition and nutritional status, influence the production of secondary metabolites in higher plants⁷ The seasonal variations in the production of phenolic compounds have been reported for variety of plants, such as *Pteridium aquilinum*, *Robinsonia evenia*, and *Menziesia ferruginea*⁸. Literature survey revealed active constituent especially in steviosides in *stevia* is greatly depends on the package of practices for the cultivation of *Stevia* and adoption of modern agro- techniques⁹. Present study concluded that climatic and habitat conditions of Uttarakhand region favorable for production of such glycosides (i.e. stevioside, rebaudioside). This is needed to warrant for detail investigations. also revealed that such difference of active ingredients due to environmental factors as well various physiological responses like temperature vary with altitudes edaphic features phenological of

species etc. Thus the glycoside contents influence probably more due to environmental factors. It is further recommended that such locations planting material used for mass multiplications for production of valuable glycoside

ACKNOWLEDGMENT

Authors thankful to Prof. A.N. Purohit, Padmashri Former M.L. Bhartia Chair for his encouragement and valuable guidance. We also thank colleagues of centre for support and help.

REFERENCES

1. Megeji NW, Kumar JK, Singh V, Kaul VK, Ahuja PS. Introducing *Stevia rebaudiana* a natural zero- caloric sweetener, Current Science 2005; 88: 801-804.
2. Kinghron AD. Biologically Active Compounds from plants with reputed medical sweeteing Properties, J. Nat. Prod, 1987; 50: 1009- 1024.

3. Stevia Facts : Natural stevia plant extract weep page on the internet retrained on 1/02/08 cited 9.2.08 available from http% presence om out text 11122341279234 – 2458/cit. Stevia – facts.
4. Mourey, D. 1992: Life with Stevia : How sweet it is OP.CIT-PP 9.
5. Zhang SQ, Kutowy O, Kumar A. *Stevia rebaudiana* leaves –A cone caloric source of sweetners, *Actulitechemiue canadienne*, 1999; 5: 22-23.
6. Chauhan N, Lohani H, Sah S, Yadav RK. *Stevai* Cultivation Techniques HRDI Centre for Aromatic Plants Dehradun India, 2005. ISBN: 81-89717-05-07.
7. Harborne JB. Introduction of Ecological Biochemistry, Academic Press, London 1988.
8. Cooper – Driver G, Finch S, Swain T. Seasonal variation in secondary plant compounds in relation to Palatability of *Pteridium aquilnum*. Biochemical and systematic Ecology, 1977; 5: 177-183.
9. Genus, J. M. C. Molecules of interest Stevioside, Phytochemistry 2003; 6: 913-921.

Table-1, Percentage of stevioside and rebaudioside Cultivated in North India

S.N.	Locality	Stevioside (%)	rebaudioside (%)
1.	Vikasnagar	9.5	3.90
2.	Sahaspur	8.82	3.30
3.	Selaqui	8.37	2.17
4.	Herbatpur	8.30	2.07
5.	Jaspur	7.40	3.20
6.	Haldwani	8.06	1.60
7.	Pantnagar	7.60	2.30
8.	Meeruth	7.11	1.80
9.	Panjab	6.53	2.50
10.	Lucknow	6.05	3.80

Source of support: Nil, Conflict of interest: None Declared